


Alt-R CRISPR-Cas9 System:

In vitro cleavage of target DNA
with ribonucleoprotein complex



See what more we can do for you at www.idtdna.com.





Revision history

Version	Date released	Description of changes
2.2	August 2019	Corrected a component amount needed to create the RNP complex.
2.1	April 2019	Adjusted component amount needed to perform the <i>in vitro</i> digestion reaction from 100 nM to 50 nM DNA substrate.
2	July 2018	Added instructions for using Alt-R CRISPR-Cas9 sgRNA. Updated names and catalog numbers for Alt-R enzymes (V3).
1.1	May 2018	Added note about use of improved Alt-R enzymes (V3): direct substitution in protocol of V3 enzymes for original enzymes (3NLS).
1	November 2017	Original protocol



Table of contents

Revision history	2
Introduction	4
Consumables	5
Protocol	6
Prepare the double-stranded DNA template as cleavage substrate	6
Prepare the guide RNA	7
Create the RNP complex	7
Perform the <i>in vitro</i> digestion reaction	8
Visualize cleaved products	8



Introduction

This protocol describes how to use a Cas9 ribonucleoprotein (RNP) complex to enable *in vitro* cleavage of double-stranded, targeted DNA. The Cas9 RNP complex contains both an Alt-R CRISPR-Cas9 guide RNA (crRNA:tracrRNA duplex or sgRNA) and an *S. pyogenes* Cas9 endonuclease. This protocol demonstrates a method to experimentally validate the activity of CRISPR guide RNA before practical application.



Consumables

Reagents	Ordering information
Option 1, 2-part guide RNA (crRNA + tracrRNA):	
<ul style="list-style-type: none">Alt-R CRISPR-Cas9 crRNA or Alt-R CRISPR-Cas9 crRNA XTAlt-R CRISPR-Cas9 tracrRNA or Alt-R CRISPR-Cas9 tracrRNA – ATTO™ 550	IDT predesigned and custom crRNA (www.idtdna.com/CRISPR-Cas9) IDT (cat # 1072532, 1072533, 1072534) IDT (cat # 1075927, 1075928)
Option 2, single guide RNA (sgRNA): Alt-R CRISPR-Cas9 sgRNA	IDT predesigned and custom sgRNA: (www.idtdna.com/CRISPR-Cas9)
Alt-R S.p. Cas9 Nuclease V3	IDT (cat # 1081058, 1081059)
Alternative: Alt-R S.p. HiFi Cas9 Nuclease V3	IDT (cat # 1081060, 1081061)
Nuclease-Free Duplex Buffer	IDT (cat # 11-05-01-12, various sizes available)
Nuclease-Free IDTE, pH 7.5 (1X TE solution)	IDT (cat # 11-01-02-02)
Nuclease-Free Water	IDT (cat # 11-04-02-01)
DNA substrate containing the target sequence	gBlocks Gene Fragments (www.idtdna.com/gBlocks), or similar
For 10X Cas9 Nuclease Reaction Buffer combine:	
<ul style="list-style-type: none">200 mM HEPES1 M NaCl50 mM MgCl₂1 mM EDTA, pH 6.5 at 25°C	General laboratory supplier
PBS	
Alternative: For Cas9 Dilution Buffer, combine:	General laboratory supplier
<ul style="list-style-type: none">30 mM HEPES150 mM KCl, pH 7.5	
Proteinase K (Molecular biology grade)	General laboratory supplier



Protocol

Prepare the double-stranded DNA template as cleavage substrate

Design your template, considering the following:

- Multiple types of double-stranded DNA can be used as substrates for Cas9 cleavage. Three common examples:
 - linearized plasmid
 - purified PCR products
 - duplexed synthetic oligos
- Your template must contain a 20 nt guide sequence, followed by the Cas9 PAM site (NGG).
- The guide sequence should match the target-specific guide RNA that will be used in the **Perform the in vitro digestion reaction**.
- The amount of template needed for the digestion may vary depending on the detection method used to **Visualize cleaved products** and the template size, as shown in this table:

DNA template	Length (bp)	Final concentration	Visualization method
DNA oligo duplex	30–100	2–5 μ M	PAGE
gBlocks fragment or PCR product	100–500	5–50 nM	Fragment Analyzer, agarose gel
gBlocks fragment or PCR product	500–2000	2–5 nM	Fragment Analyzer, agarose gel
Linearized plasmid	\geq 2000	1–2 nM	Fragment Analyzer, agarose gel

1. Ensure you are using a 10:1 molar ratio of Cas9 RNP:DNA substrate to obtain the best cleavage efficiency.
2. Resuspend or dilute the DNA substrate in Nuclease-Free Water to the required concentration.



Note: You can use the IDT resuspension calculator at www.idtdna.com/SciTools.

Prepare the guide RNA

1. Resuspend each RNA oligo (Alt-R CRISPR-Cas9 crRNA, tracrRNA, sgRNA) in IDTE buffer to a final concentration of 100 μM .
2. If you are using sgRNA, go to the next section: **Create the RNP complex**.
3. Mix the crRNA and tracrRNA oligos in equimolar concentrations in a sterile microcentrifuge tube to a final duplex concentration of 10 μM . The following table shows an example of a 10 μL final volume duplex:

Component	Amount (μL)
100 μM Alt-R CRISPR-Cas9 crRNA	1
100 μM Alt-R CRISPR-Cas9 tracrRNA	1
Nuclease-Free Duplex Buffer	8
Total volume	10

4. Heat the duplex at 95°C for 5 min.
5. Remove from heat and allow to cool to room temperature (15–25°C).

Create the RNP complex

1. Combine the guide RNA and Cas9 enzyme in equimolar amounts.

Component	Amount (μL)
100 μM Alt-R guide RNA [From Prepare the guide RNA , step 2 (sgRNA) or step 5 (crRNA:tracrRNA)]	10
Alt-R S.p. Cas9 enzyme (62 μM stock)*	1.6
PBS†	88.4
Total volume	100

* All Alt-R S.p. Cas9 enzymes are provided at a stock concentration of 62 μM .

† Cas9 RNP complexes can be made in PBS or in Cas9 dilution buffer (30 mM HEPES, 150 mM KCl, pH 7.5).

2. Incubate at room temperature for 5–10 min for optimal formation of the RNP complex.

Perform the *in vitro* digestion reaction

1. Assemble the reaction at room temperature (15–25°C).

Component	Amount (μL)
10X Cas9 Nuclease Reaction Buffer	1
1 μM Cas9 RNP	1
50 nM DNA substrate	1
Nuclease-Free Water	7
Total volume	10

2. Incubate the reaction at 37°C for 60 min.
3. Add 1 μL Proteinase K (20 mg/mL) to the reaction, then incubate the mixture at 56°C for 10 min to release the DNA substrate from the Cas9 endonuclease.

Visualize cleaved products

Analyze the digestion by using one of the following methods:

- Agarose gel electrophoresis
- Fragment Analyzer™ System (Advanced Analytical), or similar




Figure 1. Sample data showing *in vitro* digestion reaction using Alt-R tracrRNA with Alt-R CRISPR-Cas9 Positive Control crRNA (HPRT). Column-purified PCR product consisting of the Hs *HPRT* crRNA positive control sequence was used as a template in a 10 μL *in vitro* Cas9 digestion reaction. Sample 1 contains template without RNP. Sample 2 contains template and RNP. Digestion reactions were analyzed on a Fragment Analyzer system and a gel imaging system. Trace (left) shows results from Sample 2. Gel image (right) shows results for Samples 1 and 2.

Alt-R CRISPR-Cas9 System:

In vitro cleavage of target DNA with RNP complex

Integrated DNA Technologies, Inc. (IDT) is your Advocate for the Genomics Age. For more than 30 years, IDT's innovative tools and solutions for genomics applications have been driving advances that inspire scientists to dream big and achieve their next breakthroughs. IDT develops, manufactures, and markets nucleic acid products that support the life sciences industry in the areas of academic and commercial research, agriculture, medical diagnostics, and pharmaceutical development. We have a global reach with personalized customer service. See what more we can do for **you** at www.idtdna.com.

Technical support:
applicationsupport@idtdna.com

For Research Use Only. Not for use in diagnostic procedures.

© 2019 Integrated DNA Technologies, Inc. All rights reserved. Alt-R and gBlocks are trademarks of Integrated DNA Technologies, Inc., and are registered in the USA. ATTO is a trademark of ATTO-TEC GmbH. All other marks are the property of their respective owners. For specific trademark and licensing information, see www.idtdna.com/trademarks.
CRS-10096-PR 08/2019