

# HOMOLOGY-DIRECTED REPAIR USING THE Alt-R™ CRISPR-Cas9 SYSTEM AND MEGAMER™ ssDNA FRAGMENTS

## Simultaneous delivery of RNP complexes and ssDNA repair templates using the Neon™ Transfection System

### Prepare CRISPR reagents

Resuspend your oligos in Nuclease-Free IDTE, pH 7.5.

Component	Final concentration
Alt-R crRNA with tracrRNA or Alt-R sgRNA	100 $\mu$ M
Alt-R Cas9 Electroporation Enhancer	100 $\mu$ M
Megamer ssDNA fragment	0.5 $\mu$ g/ $\mu$ L, or an optimal concentration for your planned experiment (See <b>Tips</b> below)

**Note:** For assistance, use the IDT Resuspension Calculator at [www.idtdna.com/calc/resuspension](http://www.idtdna.com/calc/resuspension).

#### Tips:

- Always store CRISPR reagents at  $-20^{\circ}\text{C}$ .
- When preparing your Megamer donor, dilute donor in Nuclease-Free IDTE, or water, so that your desired dose is delivered in a 4  $\mu$ L volume.
- The Megamer donor dose may need to be optimized for your cell type. (See the protocol [Homology-directed repair using the Alt-R CRISPR-Cas9 System and Megamer ssDNA Fragments](#)).

### Prepare the gRNA complex

**Note:** If you are preparing an sgRNA, no annealing step is required. Simply dilute the sgRNA to the desired concentration in Nuclease-Free IDTE.

Prepare a two-part gRNA complex (combining crRNA and tracrRNA), which anneals the oligos to form a guide complex.

1. Combine the following components to make the gRNA complex at a final concentration of 50  $\mu$ M.

Component	Amount ( $\mu$ L)
100 $\mu$ M Alt-R CRISPR-Cas9 crRNA	5
100 $\mu$ M Alt-R CRISPR-Cas9 tracrRNA	5
IDT Duplex Buffer (to final volume)	As needed
<b>Total volume</b>	<b>10</b>

2. Heat the mixture at  $95^{\circ}\text{C}$  for 5 minutes.
3. Cool to room temperature ( $15$ – $25^{\circ}\text{C}$ ) on the bench top.

**Stopping point (optional):** gRNA complexes can be stored at  $-20^{\circ}\text{C}$  up to 1 year.

### Prepare the RNP complex

Combining the gRNA and Cas9 Nuclease allows an RNP complex to form. Prepare the delivery mix to yield a Cas9:gRNA RNP final concentration of 2:2.4  $\mu$ M.

**Note:** You can optimize the final RNP concentration for each guide. In general, a 1–4  $\mu$ M RNP concentration allows for maximal editing.

1. Combine these components per each electroporation well:

Component	Amount ( $\mu$ L)
PBS	1.8
gRNA (50 $\mu$ M)*	0.7 (36 pmol)
Alt-R Cas9 enzyme (62 $\mu$ M) <sup>†</sup>	0.5 (30 pmol)*
<b>Total volume</b>	<b>3</b>

\* If working with Cas9-GFP or Cas9-RFP, we recommend using a 1:1.2 ratio of Cas9:gRNA, instead of a 1:1 ratio.

See the [Bio-Rad Gene Pulser® Xcell™ Electroporation System protocol](#) for reference.

<sup>†</sup> This represents the concentration of the CRISPR-Cas9 enzyme; Cas9-GFP and Cas9-RFP are provided at 52  $\mu$ M (10 mg/mL). Adjust your volume accordingly.

- Incubate at room temperature for 10–20 minutes.

**Stopping point (optional):** RNP complexes can be stored at 4°C up to 1 month, or at –80°C up to 2 years in single-use aliquots.

## Prepare the cell culture media

Prepare cell culture media with and without Alt-R HDR Enhancer V2, and prewarm to 37°C for use after electroporation.

- Prewarm 190 µL of cell culture media per electroporation sample.

**Note:** Cells will be added to this media for recovery after electroporation.

- If using enhancer, mix the Alt-R HDR Enhancer V2 with cell culture media, then aliquot in a 96-well plate.

**Tip:** To reduce sample-to-sample variability, make a stock solution of cell culture media with Alt-R HDR Enhancer V2, then aliquot to the final culture plate.

- Prewarm the plate in a tissue culture incubator.

This media will be used for culturing cells for 12–24 hr after electroporation. If desired, plate triplicate wells for each electroporation sample. The following table demonstrates the amounts needed to prepare each working solution for a final concentration of 1.0 µM Alt-R HDR Enhancer V2. Each well will require 175 µL after the electroporated cells are added. If needed, scale up according to the number of samples you have.

Component	Sample (µL)	Negative control, no enhancer (µL)	Negative control, DMSO only (µL)
0.69 mM Alt-R HDR Enhancer V2	1.7*	—	—
DMSO	—	—	1.7
Cell culture media	1000	1000	1000
<b>Total volume</b>	<b>~1000</b>	<b>1000</b>	<b>1000</b>

\* The final concentration of the Alt-R HDR Enhancer V2 may need to be optimized for your cell type (see [Minimize cytotoxicity when using Alt-R HDR Enhancer V2](#)).

## Transfect cells

Prepare cells as you would for a standard CRISPR-Cas9 electroporation experiment. Make sure the cells are washed with PBS before electroporation to remove any residual nucleases.

- Suspend cells in 6 µL of the required Electroporation Buffer (e.g., R or T).
- Dilute the Alt-R Cas9 Electroporation Enhancer to a final concentration of 15 µM (if applicable).
- Make the final transfection mix by combining the following components:

Component	Amount (µL)
RNP complex	3
Megamer ssDNA donor	4
15 µM Alt-R Cas9 Electroporation Enhancer or Electroporation Buffer*	2
Cell suspension	6
<b>Total volume</b>	<b>15</b>

\* The final concentration of Alt-R Cas9 Electroporation Enhancer may need to be optimized for your cell type (see [Minimize cytotoxicity when using Alt-R HDR Enhancer V2](#)).

- After mixing the transfection mix, gently pipette 10 µL into a Neon Tip that has been inserted into the Neon Pipette, being careful to avoid air bubbles.
- Transfect cells according to the manufacturer's specifications.
- After electroporation, transfer cells to wells containing 190 µL of prewarmed culture media (without Alt-R HDR Enhancer V2) per well and gently resuspend cells.
- Transfer 50 µL of resuspended cells to the culture plates containing the prewarmed 150 µL of culture media containing Alt-R HDR Enhancer V2 (if applicable).
- Incubate cells in a tissue culture incubator.

## Change media

After 12–24 hours, remove the media from the cells, and replace with fresh media without Alt-R HDR Enhancer V2.

## Isolate gDNA

You can perform genomic DNA isolation and detect mutations 48–72 hours after electroporation.

**Stopping point (optional):** gDNA can be stored for future use by following the recommendations of the gDNA isolation protocol used.

See [Homology-directed repair using the Alt-R CRISPR-Cas9 System and Megamer ssDNA Fragments](#) for the comprehensive protocol.

## Technical support: [applicationsupport@idtdna.com](mailto:applicationsupport@idtdna.com)

For more than 30 years, IDT's innovative tools and solutions for genomics applications have been driving advances that inspire scientists to dream big and achieve their next breakthroughs. IDT develops, manufactures, and markets nucleic acid products that support the life sciences industry in the areas of academic and commercial research, agriculture, medical diagnostics, and pharmaceutical development. We have a global reach with personalized customer service.

> SEE WHAT MORE WE CAN DO FOR YOU AT [WWW.IDTDNA.COM](http://WWW.IDTDNA.COM).

**For Research Use Only. Not for use in diagnostic procedures.** Unless otherwise agreed to in writing, IDT does not intend these products to be used in clinical applications and does not warrant their fitness or suitability for any clinical diagnostic use. Purchaser is solely responsible for all decisions regarding the use of these products and any associated regulatory or legal obligations.

© 2022 Integrated DNA Technologies, Inc. All rights reserved. Alt-R and Megamer are trademarks of Integrated DNA Technologies, Inc., and are registered in the USA. Neon is a trademark of Thermo Fisher Scientific. All other marks are the property of their respective owners. For specific trademark and licensing information, see [www.idtdna.com/trademarks](http://www.idtdna.com/trademarks). Doc ID: RUO21-0211\_002 02/22